

NIFS Data Reduction


Richard McDermid

South American Gemini Data Workshop

São José dos Campos, Brazil,


October 27-30, 2011

IFU Zoo: How to map 3D on 2D



IFU Techniques: Image Slicer

MIRI -
JWST




Pros:

- Compact design
- High throughput
- Easy cryogenics


Cons:

- Difficult to manufacture



Rectangular Pixels

- NIFS has different (x,y) spatial sampling
- ***Along the slice*** is sampled by the ***detector***
- ***Across the slice*** is sampled by the ***slicer***
- Cross-slice sets spectral PSF - should be sampled on ~2 pixels
- Gives rectangular spaxels on the sky




NIFS

- Near-infrared Integral Field Spectrograph
- Cryogenic slicer design
- Z,J,H,K bands, $R \sim 5,000$
- One spatial setting:
 - 3"x3" FoV
 - 0.1"x0.04" sampling
- Optimized for use with AO
- Science: young stars, exo-planets, solar system, black holes, jets, stellar populations, hi-z galaxies....

Typical NIFS Observation

- 'Before' telluric star
 - NGS-AO
 - Acquire star
 - Sequence of on/off exposures
 - Same instrument config as science (inc. e.g. field lens for LGS)
- Science observation
 - Acquisition
 - Observation sequence:
 - Arc (grating position is not 100% repeatable)
 - Sequence of on/off exposures
- 'After' telluric (if science $>\sim 1.5$ hr)
- Daytime calibrations:
 - Baseline set:
 - Flat-lamps (with darks)
 - 'Ronchi mask' flats (with dark)
 - Darks for the arc
 - Darks for science (if sky emission to be used for wavelength calibration)

Typical NIFS Data



Arranging your files - suggestion

Daycals/ - All baseline daytime calibrations

└─ YYYYMMDD/ - cals from different dates

Science/ - All science data

└─ Obj1/ - First science object

└─ └─ YYYYMMDD/ - First obs date (if split over >1 nights)

└─ └─ └─ Config/ - e.g. 'K' (if using multiple configs)

└─ └─ └─ └─ Telluric/ - telluric data for this science obs

└─ └─ └─ └─ └─ Merged/ - Merged science and subsequent analysis

Scripts/

NIFS Reduction: Example scripts


- Three IRAF scripts on the web:
 - Calibrations
 - Telluric
 - Science
- Form the basis of this tutorial
- Data set:
 - Science object (star)
 - Telluric correction star
 - Daytime calibrations
- Update the path and file numbers at the top of each script
- Excellent starting point for basic reduction

Lamp Calibrations

- Three basic calibrations:
 - Flat (DAYCAL)
 - Correct for transmission and illumination
 - Locate the spectra on the detector
 - Ronchi Mask (DAYCAL)
 - Spatial distortion
 - Arc (NIGHTCAL)
 - Wavelength calibration
- Each has associated dark frames
- May have multiple exposures to co-add
- DAYCAL are approx. 1 per observation date
- NIGHTCAL are usually once per science target, but can be common between targets if grating config not changed

Calibration 1: Flat-Field

- Step 1: Locate the spectra
 - Mask Definition File (MDF) provides relative location of slices on detector
 - Use `nfprepare` to match this to the absolute position for your data:



- Offset is stored in a new image
- This exposure is then referenced in subsequent steps that need to know where the spectra are on the chip

Calibration 1: Flat-Field

- Step 2.1: Update flat images with offset value
- Step 2.2: Generate variance and data quality extensions
- Nfprepare is called again (once) to do both these tasks:

Input file list

Reference image with shift

```
nfprepare("@flatlist", rawpath=raw_data, shiftim="s"//calflat,  
          fl_var dq+, fl_int+, fl_corr-, fl_nonl-)
```


Create variance and
data quality planes

Run interactively

- Apply same process to dark frames

Calibration 1: Flat-Field

- Step 2.3: Combine flats and darks using gemcombine:



- Repeat for darks...
- Now have 2D images with DQ and VAR extensions. Ready to go to 3D...

Calibration 1: Flat-Field

- Step 3.1: Extract the slices using nsreduce:

'cut' out the slices from the 2D image

Apply first order wavelength coordinate system

```
nsreduce("gn"//calflat, fl_nscut+, fl_nsappw+, fl_vardq+,  
        fl_sky-, fl_dark-, fl_flat-, logfile="nifs.log")
```

- Step 3.2: Create slice-by-slice flat field using nsflat:

```
nsflat("rgn"//calflat, darks="rgn"//flatdark,  
       flatfile="rn"//calflat//"_sflat", darkfile="rn"//flatdark//"_dark",  
       fl_save_dark+, process="fit", thr_flo=0.15, thr_fup=1.55,  
       fl_vardq+, logfile="nifs.log")
```

Output flat image

Lower and Upper limits for 'bad' pixels


- Divides each spectrum (row) in a slice by a fit to the average slice spectrum, with coarse renormalizing
- Also creates a bad pixel mask from the darks

Calibration 1: Flat-Field

NOAO/IRAF V2.14.1 rmcdermi@teracles.local Tue 17:58:00 20-Jul-2010

func=spline3 order=20 low_noi=3 high_noi=3 niterate=3 grow=0

total MS= 219.



Calibration 1: Flat-Field

- Step 3.3: Renormalize the slices to account for slice-to-slice variations using `nsslitfunction`:

```
nsslitfunction("rgn"//calflat, "rn"//calflat//"_flat",  
flat="rn"//calflat//"_sflat", dark="rn"//flatdark//"_dark",  
combine="median", order=3, fl_vary-, logfile="nifs.log")
```


Final flat-field correction frame

Method to collapse
in spectral
direction

Order of fit across slices


- Fits a function in spatial direction to set slice normalization
- Outputs the final flat field, with both spatial and spectral flat information

Calibration 1: Flat-Field



Calibration 1: Flat-Field

```
NOAO/IRAF V2.14.1 rmcdermi@teracles.local Tue 18:07:42 20-Jul-2010  
func=spline3, order=3, low_rej=2, high_rej=2, niterate=3, grow=0  
total=69, sample=69, rejected=4, deleted=0, RMS= 71.02  
Determine illumination interactively for tmpsflat7480toc[SCI,2] at bin  
GCALflat
```




Calibration 2: Wavelength Calibration

- Step 1: Repeat `nprepare`, `gemcombine` and `nsreduce` -> extracted slices
- Step 2: Correctly identify the arc lines, and determine the dispersion function for each slice
 - Should run this interactively the first time through to ensure correct identification of lines and appropriate fit function
 - First solution is starting point for subsequent fits
 - Should robustly determine good solution for subsequent spectra
- Result is a series of files in a 'database/' directory containing the wavelength solutions of each slice


```
nswavelength("rgn">//arc, coordli=clist, nsum=10,  
thresho=my_thresh, trace=yes, fwidth=2.0, match=-6, cradius=8.0,  
fl_inter+, nfound=10, nlost=10, logfile="nifs.log")
```

Calibration 2: Wavelength Calibration

NOAO/IRAF V2.14.1 rmcdermi@teracles.local Tue 19:39:28 20-Jul-2010
func=chebyshev, order=4, low_rej=3, high_rej=3, niterate=10, grow=0
total=29, sample=29, rejected=3, deleted=0, RMS= 0.1087




Calibration 2: Wavelength Calibration




Calibration 3: Spatial Distortion

- Need to correct for distortions along the slices, and registration between slices
- This is done using the Ronchi mask as a reference
- Analogous to wavelength calibration, but in spatial domain

NIFS: Ronchi Mask



NIFS: Ronchi Mask



Transformation to make lines straight gives geometric correction

Calibration 3: Spatial Distortion

- Step 1: Repeat `nfpprepare`, `gemcombine` and `nsreduce` -> extracted slices
- Step 2: run `nfsdist`
 - Reference peaks are very regular, so easy to fall foul of aliasing when run automatically
 - Recommend running interactively for each daycal set


```
nfsdist("rgn"//ronchiflat, fwidth=6.0, cradius=8.0, glshift=2.8,  
minsep=6.5, thresh=2000.0, nlost=3, fl_int+, logfile="nifs.log")
```

- TIP: apply the distortion correction to the Ronchi frame itself, and check its OK


Calibration 3: Spatial Distortion

NOAO/IRAF V2.14.1 rmcdermi@teracles.local Tue 19:51:17 20-Jul-2010

identify rgnN20100410S0375[SCI,1][1024,*]



Calibration 3: Spatial Distortion



Lamp Calibrations: Summary

You now have:

1. Shift reference file: "s"+calflat
2. Flat field: "rn"+calflat+"_flat"
3. Flat BPM (for DQ plane generation):
"rn"+calflat+"_flat_bpm.pl"
4. Wavelength referenced Arc: "wrn"+arc
5. Spatially referenced Ronchi Flat: "rn"+ronchiflat

Notes:

- 1-3 are files that you need
- 4 & 5 are files with associated files in the 'database/' dir
- Arcs are likely together with science data

Telluric Star

- Similar to science reduction up to a point:
 - Sky subtraction
 - Spectra extraction => 3D
 - Wavelength calibration
 - Flat fielding
- Then extract 1D spectra, co-add separate observations, and derive the telluric correction spectrum

Telluric Star

- Preliminaries:
 - Copy the calibration files you will need into telluric directory:
 - Shift file
 - Flat
 - Bad pixel mask (BPM)
 - Ronchi mask + database dir+files
 - Arc file + database dir+files
 - Make two files listing filenames with ('object') and without ('sky') star in field

Telluric Star

- Step 1.1: Run `nfpprepare`, making use of the shift file and BPM
- Step 1.2: Combine the blank sky frames:
 - Skies are close in time
 - Use `gemcombine` and your list of sky frames to create a median sky
- Step 1.3: Subtract the combined sky from each object frame with `gemarith`

Telluric Star

- Step 2.1: Run nsreduce, this time including the flat:

```
nsreduce("sn@telluriclist",outpref="r",  
flatim=cal_data// "rn" // calflat// "_flat", fl_nscut+, fl_nsappw-,  
fl_vardq+, fl_sky-, fl_dark-, fl_flat+, logfile=log_file)
```

- Step 2.2: Replace bad pixels with values interpolated from fitting neighbours

```
nffixbad("rsn@telluriclist",outpref="b",logfile=log_file)
```

– Uses the Data Quality (QD) plane

Telluric Star

- Step 3.1: Derive the 2D spectral and spatial transformation for each slice using nsfitcoords
 - This combines the ‘1D’ dispersion and distortion solutions derived separately from ns wavelength and nsdist into a 2D surface that is linear in wavelength and angular scales
 - The parameters of the fitted surface are associated to the object frame via files in the database directory


```
nsfitcoords("brsn@telluriclist", outpref="f", fl_int+,  
lamptr="wrgn">//arc, sdisttr="rgn">//ronchiflat, lxorder=3,  
lyorder=3, sxorder=3, syorder=3, logfile=log_file)
```

Nsfitcoords - spectral

NOAO/IRAF V2.14.1 rmcdermi@teracles.local Tue 22:10:43 20-Jul-2010

Function = chebyshev, xorder = 2, yorder = 2, rms = 1.594

Fit User Coordinates to Image Coordinates for fbrsnN20100401S0146_SCI_1_




Nsfitcoords - spectral

NOAO/IRAF V2.14.1 rmcdermi@teracles.local Tue 22:43:07 20-Jul-2010

Function = chebyshev, xorder = 3, yorder = 2, rms = 0.3032

Fit User Coordinates to Image Coordinates for fbrsnN20100401S0146_SCI_3_




Nsfitcoords - spatial

NOAO/IRAF V2.14.1 rmcdermi@teracles.local Tue 22:21:19 20-Jul-2010

Function = chebyshev, xorder = 2, yorder = 2, rms = 0.06883

Fit User Coordinates to Image Coordinates for fbrsnN20100401S0146_SCI_1_




Nsfitcoords - spatial

NDAO/IRAF V2.14.1 rmcdermi@teracles.local Tue 22:48:04 20-Jul-2010

Function = chebyshev, xorder = 3, yorder = 2, rms = 0.05814

Fit User Coordinates to Image Coordinates for fbrsnN20100401S0146_SCI_3_




Telluric Star

- Step 3.2: Transform the slice images to the linear physical coordinates using `nstransform`
 - Uses transforms defined by `nsfitcoords`
 - Generates slices that are sampled in constant steps of wavelength and arcsec
- This is essentially a data-cube (even though its not a cube...)
 - Can run analysis directly from this point



When is a cube not a cube?

- 'Cube' = (x, y, λ)
- Does not have to be a rectilinear 3D array



Telluric Star

- Step 4.1: Extract 1D aperture spectra from the data cube
 - Use `nfextract` to define an aperture (radius and centre) and sum spectra within it
 - Outputs a 1D spectrum
- Step 4.2: Co-add the 1D spectra using `gemcombine`



Science Data

- Same preliminaries as telluric:
 - Copy database and arc+Ronchi files
 - Copy shift file, flat and BPM
 - Identify sky and object frames
- In addition, we make use of the 1D telluric
- Generally need to combine separate (and dithered) data-cubes


Science Data

- Initial steps:
 - Nfprepare as per telluric
 - Subtract sky using gemarith
 - Usually have one unique sky per object: ABAB
 - Can have ABA – two science share a sky
 - Nsreduce (inc. flat field)
 - Nffixbad, nsfitcoords, nstransform
- Now have data-cube with linear physical coordinates


Science Data: Telluric correction

- Telluric spectrum is not only atmosphere, but also stellar spectrum:
 - Need to account for stellar absorption features
 - AND account for black-body continuum
- Needs some 'by-hand' steps to prepare the telluric star spectrum
 - Remove strong stellar features with `splot`
 - Remove BB shape with a BB spectrum

Science Data: Telluric correction




Science Data: Telluric correction




Telluric Absorption

- Alternative approach is to fit a stellar template (Vacca et al. 2003)
- Need good template
- Can use solar-type stars, but needs careful treatment...



Science Data: Telluric correction

- Finally, run nftelluric
 - Computes the normalized correction spectrum
 - Allows for shifts and amplitude scaling
 - Divides the correction spectrum through the data



Science Data: Merging

- Now have series of data-cubes:
 - No dark current or sky (sky-subtracted)
 - Spatially and spectrally linearized
 - Bad pixels interpolated over
 - No instrumental transmission (flat-fielded)
 - No atmospheric transmission (telluric-corrected)
- Need to **combine** the data-cubes


Science Data: Merging

Two approaches:

1. Dithering by non-integer number of spaxels:

- Allows over-sampling, via 'drizzling'
- Resampling introduces correlated noise
- Good for fairly bright sources


Nifcube / gemcube method



2. Dither by integer number of spaxels

- Allows direct 'shift and add' approach
- No resampling:- better error characterisation
- Assumes accurate (sub-pixel) offsetting
- No over-sampling gain
- Suitable for 'deep-field' applications

Imcombine method



Science Data: Merging with nf/gemcube

1. Use nfcube to create the 3D arrays
 - 2D slices to 3D cube - interpolation
 - Default pixel scale is 0.05"x0.05" (arrays need square pixels..)
2. Find a reference pixel coordinate
 - These cubes are easily displayed using ds9
 - Load as an array, scroll through the slices
 - Reference point should be an easily recognizable feature, common to all the cubes
 - Can blindly use offsets, but not recommended
3. Adapt the headers to reflect the common spatial axes origin
4. Run gemcube (interpolates again)

What headers to change?

- Gemcube uses the file WCS to merge overlapping cubes
- Need to 'trick' the WCS to include your offsets
- Following list seemed to work:
 - Position of reference source measured on cube plane – different in each cube
 - CRPIX1 - Reference source (x position in pixels), e.g '29'
 - CRPIX2 - Reference source (y position in pixels), e.g. '27.5'
 - Approx decimal RA & Dec at reference point – same in all cubes
 - CRVAL1 - approx. decimal RA of target (same for all cubes)
 - CRVAL2 - approx. decimal declination of target
 - Pixel scale parameters - these are set here to be 0.05 arcsec, but in degrees
 - CD1_1 = 0.0
 - CD1_2 = 1.9E-05
 - CD2_1 = -4.7e-5
 - CD2_2 = 0.0
 - Coordinate system info
 - WAT1_001 = 'wtype=linear axtype=xi'
 - WAT2_001 = 'wtype=linear axtype=eta'
 - CTYPE1 = 'RA---TAN'
 - CTYPE2 = 'DEC--TAN'




Hack alert! This is not the best way to proceed! But lets you progress within iraf. Other ideas are welcome!

Science Data: Merging with nf/gemcube

- This approach involves (at least) one superfluous interpolation: nifcube + gemcube both interpolate
- Might be possible to use gemcube directly from pre-transformed data, but wrapper not written (TBD: works on single slices, so can be adapted)
- Nifcube step is convenient for determining reference coordinate, and allows gemcube to combine the frames
- Not ideal, but gives a way to combine your data

Science Data: Merging with imcombine

1. Convert files to 3D cubes with nfcube
2. Provide the list of input files, and an 'offsets' file of integer shifts to imcombine



```
imcombine("cat*.fits", output="Merged", offset=offsets.dat,  
logfile=log_file)
```

Science Data: Merging

Other details for merging not covered:

- For smoother images / less border effects, individual cubes should be renormalized to have common flux in overlap regions
- For optimal S/N, spectra should be weighted by the variance when combining
- Even without dithers, long observations will have sub-pixel shifts due to flexure and atmospheric refraction
- To maintain high image quality, keep a reference source common to all fields in the mosaic if possible, and use sub-pixel dithers

The End

(....or rather, just the beginning....)